

Volume contents

Volume 49 (1995)

Issue contents

- K. Manova, V. De Leon, M. Angeles, S. Kalantry, M. Giarré, L. Attisano, J. Wrana, R.F. Bachvarova**
mRNAs for activin receptors II and IIB are expressed in mouse oocytes and in the epiblast of pregastrula and gastrula stage mouse embryos 1
- J. Riese, R. Zeller, R. Dono**
Nucleo-cytoplasmic translocation and secretion of fibroblast growth factor-2 during avian gastrulation 13
- D. Uwanogho, M. Rex, E.J. Cartwright, G. Pearl, C. Healy, P.J. Scotting, P.T. Sharpe**
Embryonic expression of the chicken *Sox2*, *Sox3* and *Sox11* genes suggests an interactive role in neuronal development 23
- S. Stein, M. Kessel**
A homeobox gene involved in node, notochord and neural plate formation of chick embryos 37
- L. Bally-Cuif, M. Gulisano, V. Broccoli, E. Boncinelli**
c-otx2 is expressed in two different phases of gastrulation and is sensitive to retinoic acid treatment in chick embryo 49
- I. Minoura, H. Nakamura, K. Tashiro, K. Shiokawa**
Stimulation of circus movement by activin, bFGF and TGF- β 2 in isolated animal cap cells of *Xenopus laevis* 65
- S. Ibensouda, F. Schweisguth, D. Jullien, C. Kücherer, J.-A. Lepesant, A. Vincent**
Evolutionarily conserved positive and negative cis-acting elements control the blastoderm-specific expression of the *Drosophila serendipity* α cellularisation gene 71
- M. Mével-Ninio, R. Terracol, C. Salles, A. Vincent, F. Payre**
ovo, a *Drosophila* gene required for ovarian development, is specifically expressed in the germline and shares most of its coding sequences with *shavenbaby*, a gene involved in embryo patterning 83

- M. Haenlin, W.F. McDonald, C. Cibert, E. Mohier**
The angle of the dorsoventral axis with respect to the anteroposterior axis in the *Drosophila* embryo is controlled by the distribution of *gurken* mRNA in the oocyte 97
- S.J. Poole**
Conservation of complex expression domains of the *pdm-2* POU domain gene between *Drosophila virilis* and *Drosophila melanogaster* 107
- T.A. Rosenquist, G.R. Martin**
Visceral endoderm-1 (VE-1): an antigen marker that distinguishes anterior from posterior embryonic visceral endoderm in the early post-implantation mouse embryo 117
- H. Nakamura, K. Tashiro, T. Nakamura, K. Shiokawa**
Molecular cloning of *Xenopus* HGF cDNA and its expression studies in *Xenopus* early embryogenesis 123
- J. Reinitz, D.H. Sharp**
Mechanism of *eve* stripe formation 133
- Issue contents 159
- P.P. D'Avino, S. Crispi, L.C. Polito, M. Furia**
The role of the *BR-C* locus on the expression of genes located at the ecdysone-regulated 3C puff of *Drosophila melanogaster* 161
- N.E. Baker, A.E. Zitron**
Drosophila eye development: *Notch* and *Delta* amplify a neurogenic pattern conferred on the morphogenetic furrow by *scabrous* 173
- M.E. Lane, D. Kalderon**
Localization and functions of Protein Kinase A during *Drosophila* oogenesis 191
- F.L. Conlon, C.V.E. Wright, E.J. Robertson**
Effects of the *T^W* mutation on notochord formation and mesodermal patterning 201

R. Wilson, T. Mohun		family defines dorsal regions of the neural plate and the embryonic retina	223
XIdx, A dominant negative regulator of bHLH function in early <i>Xenopus</i> embryos	211		
M. Rissi, J. Wittbrodt, E. Délot, M. Naegeli, F.M. Rosa		Author index	235
Zebrafish <i>Radar</i> : A new member of the TGF- β super-		Subject index	237
		Volume contents	241

